



TAF1 gene

TATA-box binding protein associated factor 1

Normal Function

The *TAF1* gene provides instructions for making part of a protein called transcription factor IID (TFIID). This protein is active in cells and tissues throughout the body, where it attaches (binds) to DNA. Transcription factor IID plays an essential role in regulating the activity of most genes.

The *TAF1* gene is part of a complex region of DNA known as the *TAF1/DYT3* multiple transcript system. This region consists of short stretches of DNA from the *TAF1* gene plus some extra segments of genetic material near the gene. These stretches of DNA can be combined in different ways to create various sets of instructions for making proteins. Researchers believe that some of these variations are critical for the normal function of nerve cells (neurons) in the brain.

Health Conditions Related to Genetic Changes

X-linked dystonia-parkinsonism

Several changes in the *TAF1/DYT3* multiple transcript system have been identified in people with X-linked dystonia-parkinsonism. Some alter single DNA building blocks (nucleotides) in the gene; these changes are described as disease-specific single-nucleotide changes (DSCs). Another genetic change deletes a small number of nucleotides from the gene. Researchers are uncertain how these changes are related to the movement abnormalities characteristic of the disease.


X-linked dystonia-parkinsonism may also be related to an extra segment of DNA in the *TAF1/DYT3* multiple transcript system. The extra segment results from the insertion of a retrotransposon, which is a small piece of DNA that can move around to different positions in a cell's genetic material. When a retrotransposon inserts itself in or a near a gene, it can disrupt the gene's function. In this case, the retrotransposon insertion probably interferes with the normal function of the *TAF1/DYT3* multiple transcript system.

Researchers suspect that changes in the *TAF1/DYT3* multiple transcript system disrupt the regulation of critical genes in neurons. This defect leads to the eventual death of these cells, particularly in areas of the brain called the caudate nucleus and putamen. These regions are critical for normal movement, learning, and memory. It is unclear why the effects of changes in the *TAF1/DYT3* multiple transcript system appear to be limited to dystonia and parkinsonism.

Chromosomal Location

Cytogenetic Location: Xq13.1, which is the long (q) arm of the X chromosome at position 13.1

Molecular Location: base pairs 71,366,220 to 71,530,525 on the X chromosome (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- BA2R
- CCG1
- CCGS
- Cell cycle gene 1 protein
- cell cycle, G1 phase defect
- complementation of cell cycle block, G1-to-S
- DYT3
- DYT3/TAF1
- KAT4
- N-TAF1
- NSCL2
- P250
- TAF(II)250
- TAF1 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 250kDa
- TAF1_HUMAN
- TAF2A
- TAFII-250

- TAFII250
- TATA box-binding protein-associated factor 1
- TATA box binding protein associated factor 1
- TATA box-binding protein-associated factor 2A
- TBP-associated factor 1
- TBP-associated factor 250 kDa
- TBP-associated factor, RNA polymerase II, 250-kD
- transcription factor TFIID p250 polypeptide
- Transcription initiation factor TFIID 250 kDa subunit
- Transcription initiation factor TFIID subunit 1

Additional Information & Resources

Educational Resources

- Molecular Biology of the Cell (fourth edition, 2002): A Large Fraction of the Human Genome Is Composed of Nonretroviral Retrotransposons
<https://www.ncbi.nlm.nih.gov/books/NBK26845/#A884>
- Molecular Biology of the Cell (fourth edition, 2002): RNA Polymerase II Requires General Transcription Factors
<https://www.ncbi.nlm.nih.gov/books/NBK26887/#A999>

GeneReviews

- X-Linked Dystonia-Parkinsonism Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1489>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28TAF1%5BTIAB%5D%29+OR+%28DYT3%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- TAF1 RNA POLYMERASE II, TATA BOX-BINDING PROTEIN-ASSOCIATED FACTOR, 250-KD
<http://omim.org/entry/313650>

Research Resources

- **Atlas of Genetics and Cytogenetics in Oncology and Haematology**
http://atlasgeneticsoncology.org/Genes/GC_TAF1.html
- **ClinVar**
<https://www.ncbi.nlm.nih.gov/clinvar?term=TAF1%5Bgene%5D>
- **HGNC Gene Family: Lysine acetyltransferases**
<http://www.genenames.org/cgi-bin/genefamilies/set/486>
- **HGNC Gene Symbol Report**
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=11535
- **NCBI Gene**
<https://www.ncbi.nlm.nih.gov/gene/6872>
- **UniProt**
<http://www.uniprot.org/uniprot/P21675>

Sources for This Summary

- Herzfeld T, Nolte D, Müller U. Structural and functional analysis of the human TAF1/DYT3 multiple transcript system. *Mamm Genome.* 2007 Nov;18(11):787-95. Epub 2007 Oct 19.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17952504>
- Maile T, Kwoczynski S, Katzenberger RJ, Wassarman DA, Sauer F. TAF1 activates transcription by phosphorylation of serine 33 in histone H2B. *Science.* 2004 May 14;304(5673):1010-4. Retraction in: McNutt M. *Science.* 2014 May 30;344(6187):981.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15143281>
- Makino S, Kaji R, Ando S, Tomizawa M, Yasuno K, Goto S, Matsumoto S, Tabuena MD, Maranon E, Dantes M, Lee LV, Ogasawara K, Tooyama I, Akatsu H, Nishimura M, Tamiya G. Reduced neuron-specific expression of the TAF1 gene is associated with X-linked dystonia-parkinsonism. *Am J Hum Genet.* 2007 Mar;80(3):393-406. Epub 2007 Jan 23.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17273961>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1821114/>
- Muller U, Herzfeld T, Nolte D. The TAF1/DYT3 multiple transcript system in X-linked dystonia-parkinsonism. *Am J Hum Genet.* 2007 Aug;81(2):415-7; author reply 417-8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17668393>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1950800/>
- Nolte D, Niemann S, Müller U. Specific sequence changes in multiple transcript system DYT3 are associated with X-linked dystonia parkinsonism. *Proc Natl Acad Sci U S A.* 2003 Sep 2;100(18):10347-52. Epub 2003 Aug 19.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12928496>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC193564/>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/TAF1>

Reviewed: December 2008

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services